



Burden of obesity in the Eastern Mediterranean Region: findings from the Global Burden of Disease 2015 study

GBD 2015 Eastern Mediterranean Region Obesity Collaborators¹

Received: 1 May 2017/Revised: 21 June 2017/Accepted: 23 June 2017/Published online: 3 August 2017
© The Author(s) 2017. This article is an open access publication

Abstract

Objectives We used the Global Burden of Disease (GBD) 2015 study results to explore the burden of high body mass index (BMI) in the Eastern Mediterranean Region (EMR). **Methods** We estimated the prevalence of overweight and obesity among children (2–19 years) and adults (≥ 20 years) in 1980 and 2015. The burden of disease related to high BMI was calculated using the GBD comparative risk assessment approach.

Results The prevalence of obesity increased for adults from 15.1% (95% UI 13.4–16.9) in 1980 to 20.7% (95% UI 18.8–22.8) in 2015. It increased from 4.1% (95% UI 2.9–5.5) to 4.9% (95% UI 3.6–6.4) for the same period among children. In 2015, there were 417,115 deaths and 14,448,548 disability-adjusted life years (DALYs) attributable to high BMI in EMR, which constitute about 10 and 6.3% of total deaths and DALYs, respectively, for all ages.

Conclusions This is the first study to estimate trends in obesity burden for the EMR from 1980 to 2015. We call for EMR countries to invest more resources in prevention and health promotion efforts to reduce this burden.

Keywords Obesity · Burden of disease · Eastern Mediterranean Region

Introduction

High body mass index (BMI), or overweight, is associated with increased morbidity and mortality, and is a major risk factor for diabetes, cancer, cardiovascular diseases, sleep apnea, and poor physical health (Kim et al. 2016; Yao et al. 2017; Kelly et al. 2017; Mehta et al. 2017). It is also associated with an increased risk for psychiatric disorders, including depression (Pratt and Brody 2014; Abou Abbas et al. 2015). A continuous rise in obesity is threatening health improvements in many countries (Sidney et al. 2016; GBD 2015 Obesity Collaborators 2017), while controlling its spread could drastically improve population health (Maciosek et al. 2017).

The increase in overweight and obesity prevalence is a direct result of lifestyle changes due to the social and demographic transition that started several decades ago (Broyles et al. 2015; Mokdad et al. 2016). The Eastern Mediterranean Region (EMR) is facing the same challenges due to rapid economic, demographic, and lifestyle changes, including changes in food consumption, reduced physical activity, and increased sedentary lifestyle (Musaiger et al. 2012; Mokdad et al. 2016). The contribution of high BMI to total DALYs in the EMR had increased in 2013 (7.5% of DALYs) in comparison to 1990 (3.7% of DALYs) (Mokdad et al. 2016). Kuwait, Qatar, and Libya,

This article is part of the supplement “The state of health in the Eastern Mediterranean Region, 1990–2015.”

The members of GBD (Global Burden of Disease) 2015 Eastern Mediterranean Region Obesity Collaborators are listed at the end of the article. Ali H. Mokdad, on behalf of GBD 2015 Eastern Mediterranean Region Obesity Collaborators, is the corresponding author.

Electronic supplementary material The online version of this article (doi:10.1007/s00038-017-1002-5) contains supplementary material, which is available to authorized users.

✉ GBD 2015 Eastern Mediterranean Region Obesity Collaborators
mokdaa@uw.edu

¹ Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, USA

three EMR countries, were among the top ten countries with highest prevalence of obesity worldwide in 2013, and the man believed to be the heaviest living person was diagnosed in Saudi Arabia (Ng et al. 2014; Terkawi et al. 2014).

The EMR has a population of about 583 million people (World Health Organization 2016). Countries in the EMR vary significantly in terms of their gross domestic product, socio-demographic profiles, health indicators, and health system capacities and coverage. Despite the heavy burden of high BMI in the region, no comprehensive and current estimates of the epidemic exist for the EMR.

To quantify the burden of high BMI in the EMR and its impact on health, we systematically evaluated the trends in prevalence of overweight and obesity as well as the patterns of deaths and DALYs related to high BMI by age and sex, using the results of the Global Burden of Disease (GBD) 2015 study. We also estimated the attributable fraction of high BMI to ischemic heart disease, stroke, and diabetes mellitus, the three leading non-communicable causes of death in EMR, for which high BMI is a risk factor.

Methods

The prevalence of overweight and obesity among children (2–19 years) and adults (≥ 20 years) in 1980 and 2015 was estimated for EMR countries. The EMR countries, based on the World Health Organization classification, are Afghanistan, the Kingdom of Bahrain, Djibouti, the Arab Republic of Egypt, the Islamic Republic of Iran, the Republic of Iraq, the Hashemite Kingdom of Jordan, the State of Kuwait, the State of Lebanon, the State of Libya, the Kingdom of Morocco, the Sultanate of Oman, the Islamic Republic of Pakistan, Palestine, the State of Qatar, the Kingdom of Saudi Arabia, the Federal Republic of Somalia, the Republic of Sudan, the Syrian Arab Republic, the Republic of Tunisia, the United Arab Emirates, and the Republic of Yemen.

The burden of disease related to high BMI was calculated using the GBD comparative risk assessment approach between 1990 and 2015 (Forouzanfar et al. 2015, 2016). A detailed methodology of BMI estimation for GBD 2015 has been published elsewhere (GBD 2015 Obesity Collaborators 2017). Since burden estimations depend on GBD all-cause mortality, burden of high BMI is only available for the period 1990–2015. We used all available data surveys following a systematic search. The search strategy as well as data sources used per country have been published as an appendix elsewhere and are available from the Global Health Data Exchange (Institute for Health Metrics and Evaluation 2016; GBD 2015 Obesity Collaborators 2017).

Briefly, Medline was systematically searched for studies providing nationally or subnationally representative estimates of overweight prevalence, obesity prevalence, or mean body mass index (BMI) published between 1 January 2014 and 31 December 2015 to update the GBD 2013 systematic literature search (Ng et al. 2016).

For adults, 127 out of 2036 abstracts identified met inclusion criteria and were extracted. For children, 146 out of 971 articles identified were extracted. In total, 816 articles were included in the analysis. Additionally, the Global Health Data Exchange (GHDx) database was searched for individual-level data from major multinational survey series or country-specific surveys and identified 1026 unique sources meeting the inclusion criteria. Of the 816 articles and 1026 unique sources, all those pertaining to EMR countries were included. The GBD 2015 results tool from the GHDx allows readers to view, country by country, what data sources have been used to produce these estimates (Institute for Health Metrics and Evaluation 2016).

For adults, overweight was defined as $25.0 \leq \text{BMI} \leq 30 \text{ kg/m}^2$, and obesity was defined as $\text{BMI} \geq 30 \text{ kg/m}^2$. The International Obesity Task Force definition was used for childhood overweight and obesity (Cole et al. 2000).

Children were defined as individuals 2–19 years of age based on the lowest age for which the International Obesity Task Force provides a definition of overweight and obesity, and the age groups used in GBD modeling, which include 19 in the age group 15–19 (Cole et al. 2000; Wang et al. 2016).

Briefly, a spatiotemporal Gaussian process regression (ST-GPR) was used to estimate the mean prevalence of overweight and obesity (Ng et al. 2014). To improve estimates for countries with sparse data, three country-level covariates with best fit and coefficients in the expected direction were selected: 10-year lag-distributed energy intake per capita, the absolute latitude of the country as proxy for income, and the proportion of people living in urban areas. These covariates have been systematically evaluated in a previous study (Ng et al. 2014).

The Bradford Hill criteria for causation and the World Cancer Research Fund evidence grading criteria were used to systematically evaluate epidemiologic evidence supporting the causal relationship between high BMI and various diseases among adults (≥ 20 years of age) (Hill 1965; World Cancer Research Fund and American Institute for Cancer Research 2007).

The population-attributable fraction by country, age, sex, and year was calculated to quantify the burden of disease related to high BMI, defined as $\text{BMI} \geq 25 \text{ kg/m}^2$, for each disease. Deaths and DALYs related to high BMI for each country, age, sex, year, and cause were computed

by multiplying the population-attributable fraction by the total deaths or DALYs estimated in GBD 2015 for that country, age, sex, year, and cause. The total disease burden of high BMI was calculated as the sum of disease-specific burden.

95% uncertainty intervals (UI) for all results were computed using Monte Carlo simulations, keeping 1000 draws of each quantity of interest to propagate uncertainty into final estimates.

Expected estimates were also produced for each country based on its Socio-demographic Index (SDI)—a summary measure of lag-distributed income per capita, average educational attainment over the age of 15 years, and total fertility rate (Forouzanfar et al. 2015). In GBD 2015, SDI was computed by rescaling each component to a scale of zero to one, with zero being the lowest observed educational attainment, lowest income per capita, and highest fertility rate from 1980 to 2015, and one being the highest observed educational attainment, highest income per capita, and lowest fertility rate during that time, and then taking the geometric mean of these values for each location-year.

This study followed the Guidelines for Accurate and Transparent Health Estimates Reporting (GATHER) of the World Health Organization (WHO) regarding documentation of data sources, estimation methods, and statistical analysis (Stevens et al. 2016).

Role of the funding source

The Bill & Melinda Gates Foundation had no role in the development of these methods.

Results

The mean BMI increased from 25.2 kg/m² [95% uncertainty interval (UI) 24.9–25.5] in 1980 to 26.0 kg/m² (95% UI 25.8–26.3) in 2015 in the EMR among persons aged 20 years or older. The prevalence of obesity increased from 15.1% (95% UI 13.4–16.9) to 20.7% (95% UI 18.8–22.8) for the same period and for the same age group (Fig. 1). It increased from 4.1% (95% UI 2.9–5.5) to 4.9% (95% UI 3.6–6.4) for the same period among those aged 2–19 years (Fig. 1). The highest prevalence of obesity among adults 20 years or older in 2015 was observed in Qatar: 42.5% (95% UI 40.1–44.8) for males and 52.4% (95% UI 50.3–54.5) for females (e-Table 1); and the highest for children 2–19 years was observed in Kuwait: 22.1% (95% UI 17.8–27.0) for males and 19.2% (95% UI 15.2–23.4) for females (e-Table 2). The lowest prevalence of obesity was observed in Somalia among individuals 20 years or older: 2.5% (95% UI 1.5–4.0) for males and 10.5% (95% UI




Fig. 1 Prevalence of obesity among adults aged 20 years or older and children aged 2–19 years in 1980 and 2015 (Global Burden of Disease 2015 study, Eastern Mediterranean Countries, 1980 and 2015)

8.3–12.8) for females (e-Table 1), and in Yemen and Pakistan for children 2–19 years of age: 1.3% (95% UI 0.9–1.8) for males in Yemen and 2.2% (95% UI 1.4–3.3) for females in Pakistan (e-Table 2). Prevalence of obesity was higher in females than males 20 years or older for all countries, with Sudan having the highest difference between sexes: 11.4% (95% UI 10.0–13.1) for males and 28.3% (95% UI 25.6–31.2) for females (e-Table 1). The highest difference in obesity prevalence between sexes for children was observed in Qatar: 20.8% (95% UI 16.5–25.1) for males and 13.5% (95% UI 10.3–17.1) for females. In children, prevalence of obesity was higher in males for several countries (e-Table 2).

Deaths

In 2015, there were 417,115 deaths attributable to high BMI in EMR, which constitute about 10% of total deaths in the region for all ages. This is a rate of 120.1 (95% UI 87.5–156.2) deaths per 100,000 population, an 11% increase since 1990. It contributed to 5.0, 0.9, and 1.9% of all deaths cause by ischemic heart disease, ischemic stroke, and diabetes mellitus, respectively. Contribution by specific age groups is detailed in Table 1. In 2015, the rate of deaths attributable to high BMI was highest in Afghanistan, 227.6 (95% UI 146.2–319.5), and lowest in Tunisia, 64.8 (95% UI 42.7–92.5) per 100,000 population (Table 2). Overall, death rates attributable to high BMI have increased in 11 countries and decreased in 11 (Table 2). The largest increase in deaths per 100,000 population attributable to high BMI was observed in Djibouti: from 37.3 (95% UI 15.1–70.6) in 1990 to 101.1 (95% UI 47.5–192.7) in 2015 (Table 2). The largest decrease was observed in Lebanon: from 124.8 (95% UI 85.2–169.7) in 1990 to 72.5 (95% UI 47.0–101.0) in 2015 (Table 2).

Table 1 Deaths, with 95% uncertainty intervals (UI), per 100,000 population attributable to high body mass index among those who died from ischemic heart disease, stroke, and diabetes, by age groups

(Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990 and 2015)

Age	Cause	1990		2015	
		Males	Females	Males	Females
15–49	Ischemic heart disease	10.1 (6.0–14.7)	7.0 (4.5–9.5)	12.1 (7.6–17.2)	6.9 (4.7–9.3)
	Stroke	7.1 (4.6–9.8)	8.1 (5.8–10.6)	7.2 (4.7–9.9)	7.3 (5.3–9.6)
	Diabetes	1.7 (1.2–2.3)	2.0 (1.5–2.6)	2.6 (1.8–3.3)	2.9 (2.2–3.6)
50–69	Ischemic heart disease	118.4 (67.9–177.4)	118.1 (80.0–161.8)	134.4 (82.3–191.6)	107.1 (74.8–142.7)
	Stroke	58.1 (33.6–85.1)	77.8 (53.9–107.7)	60.6 (38.4–85.5)	68.7 (49.1–91.2)
	Diabetes	24.8 (15.7–34.4)	36.4 (25.1–50.1)	39.2 (27.2–52.3)	53.4 (40.6–67.8)
70+	Ischemic heart disease	283.8 (141.5–460.9)	366.1 (225.5–539.6)	336.9 (183.3–532.9)	361.0 (222.4–520.7)
	Stroke	117.4 (58.4–198.2)	163.6 (96.9–248.3)	125.8 (65.8–203.8)	150.7 (89.1–223.9)
	Diabetes	57.7 (30.8–89.2)	81.9 (47.6–130.2)	106.5 (59.1–163.1)	160.3 (103.1–224.8)

DALYs

In 2015, there were 14,448,548 DALYs attributable to high BMI in the EMR, which constitutes about 6.3% of total DALYs in the region for all ages. This is a rate of 3452.5 (95% UI 2599.2–4386.5) DALYs per 100,000 population, a 13.9% increase since 1990. It contributed to 3.0, 0.5, and 2.3% of all DALYs caused by ischemic heart disease, stroke, and diabetes mellitus, respectively. Contribution by specific age groups is detailed in Table 3. In 2015, the rate of DALYs attributable to high BMI was highest in Afghanistan, 6576.7 (95% UI 4366.1–9219.6), and lowest in Tunisia, 2022.4 (95% UI 1395.1–2724.3) per 100,000 population. Trends in DALYs followed trends in deaths from high BMI in all countries. The largest increase in DALYs per 100,000 population attributable to high BMI was observed in Djibouti: from 1111.0 (95% UI 492.8–2005.8) in 1990 to 2810.8 (95% UI 1432.5–5212.9) in 2015 (Table 4). The largest decrease was observed in Lebanon: from 3552.4 (95% UI 2508.2–4645.4) in 1990 to 2363.8 (95% UI 1707.9–3077.2) in 2015.

Expected versus observed

Overall, and based on an SDI of 0.55, a death rate of 74.0 and a DALYs rate of 2114.3 per 100,000 population were expected for the EMR in 2015 for high BMI. Both are lower than the observed death rate of 120.1 (95% UI 87.5–156.2) and DALYs rate of 3452.5 (95% UI 2599.2–4386.5) for the same year. Expected estimates for each country are detailed in Tables 3 and 4.

Discussion

This is the first study to provide estimates of trends in obesity prevalence, deaths, and DALYs for the EMR from 1980 to 2015. Our study showed that the observed burden

for high BMI was higher than expected for most countries in the region based on their SDI levels. Our study calls for renewed efforts to reduce the burden of obesity in the region. Indeed, with further progression of the epidemiologic transition and the growth and aging of the EMR population, high BMI will increase the burden of chronic conditions and disability and put financial and resource strains on the health systems.

High BMI is observed in some poor and rich countries of the EMR. In developed countries such as the United States and France, obesity is higher among low socioeconomic strata of the population (Drewnowski et al. 2014). These patterns in the West are attributable to poor diet and lower physical activity levels. Previous studies have reported similar findings for both socioeconomic status and sex in the EMR (Musaiger 2011a). In the EMR, except for Iran and Lebanon, all high-middle- and high-SDI countries had an obesity prevalence equaling or exceeding 25% for both males and females 20 years or older. Obesity prevalence exceeded 10% among children 2–19 years only in these countries as well, except for Bahrain, Jordan, Lebanon, and Iran. Djibouti and Egypt were the only low- and middle-SDI countries where childhood obesity exceeded 10% for females. These country-level estimates might be masking variations of the epidemic within each of the countries. Indeed, few studies have been done at the country level in the EMR, and these showed a variation in BMI levels between levels of education and income. High BMI was more likely to impact those with low educational levels (Sibai et al. 2003; Memish 2014).

Unfortunately, our estimations for observed obesity burden were higher than expected for the region, and based on SDI, which only deals with socioeconomic inequalities between countries. However, the biggest gaps between the included countries are cultural factors and obesogenic cultural traditions, as well as political instability (wars, civil unrest,) all of which are linked to obesity but not

Table 2 Deaths, with 95% uncertainty intervals (UI), per 100,000 population due to high body mass index observed in 1990 and 2015, and expected based on Socio-demographic Index in 2015 (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990 and 2015)

Country	1990 deaths		2015 deaths		Total observed		Total expected
	Males		Males		Females		
	Males	Females	Males	Females	Males	Females	
Afghanistan	166.9 (80.9–283.8)	296.3 (170.9–460.1)	160.5 (78.6–268.2)	292.1 (166.4–450.7)	27,827.4 (17,910.5–39,914.8)	5797.5	
Bahrain	133.2 (84.8–191.7)	175.2 (129.1–222.4)	97.7 (62.1–141.3)	99.2 (68.9–130.4)	557.5 (421.6–709.1)	443.3	
Djibouti	37.1 (9.8–94.0)	36.7 (11.8–82.8)	102.0 (35.3–229.3)	99.1 (37.7–231.1)	467.9 (211.4–929.9)	312.5	
Egypt	153.2 (98.2–207.3)	154.4 (116.9–192.2)	176.2 (123.4–228.0)	154.6 (121.5–188.6)	90,774.1 (73,283.6–108,972.5)	48,851.2	
Iran	90.7 (44.8–146.5)	98.9 (63.3–138.2)	102.3 (56.9–162.0)	94.4 (59.8–137.7)	49,386.0 (33,333.8–68,369.9)	44,524.9	
Iraq	234.4 (158.3–335.3)	243.7 (171.0–324.3)	217.4 (129.1–321.1)	206.3 (136.0–295.7)	30,963.7 (22,483.9–41,095.5)	12,627.7	
Jordan	144.8 (93.7–209.6)	198.5 (146.6–258.2)	118.8 (80.1–160.0)	107.3 (80.4–136.1)	3687.0 (2946.9–4478.1)	3082.9	
Kuwait	92.2 (64.5–120.0)	115.1 (89.1–142.1)	93.9 (65.2–126.1)	93.6 (70.6–120.7)	1335.8 (1071.0–1657.8)	670.7	
Lebanon	127.0 (70.4–198.5)	122.6 (76.9–179.0)	69.4 (35.3–113.1)	75.2 (44.8–109.8)	3565.9 (2320.6–4940.1)	3803.3	
Libya	81.8 (49.9–120.0)	103.5 (72.0–137.6)	96.8 (59.6–141.4)	109.7 (78.2–148.0)	3727.3 (2795.3–4725.7)	3261.4	
Morocco	85.0 (45.4–132.0)	103.3 (65.8–148.5)	77.4 (40.8–129.9)	102.6 (57.3–158.6)	22,383.7 (14,985.5–31,865.0)	18,621.5	
Oman	82.0 (46.3–131.0)	89.4 (51.9–140.9)	92.3 (56.9–131.3)	93.4 (63.0–123.8)	1575.4 (1161.4–2011.7)	1619.5	
Pakistan	54.3 (22.0–98.4)	67.1 (33.2–113.4)	105.9 (169.5–54.7)	110.6 (64.32–166.3)	110,546.2 (72,034.6–153,911.6)	67,064.8	
Palestine	114.4 (63.3–182.1)	128.0 (77.6–197.6)	149.7 (86.8–225.0)	118.2 (77.7–170.3)	2347.3 (1655.2–3168.4)	1502.5	
Qatar	142.7 (101.7–187.8)	209.5 (161.3–263.6)	106.6 (69.3–154.4)	126.1 (88.2–169.1)	567.3 (424.8–747.7)	479.1	
Saudi Arabia	75.2 (46.7–107.1)	81.9 (59.0–108.4)	84.5 (57.9–114.5)	68.3 (51.6–87.4)	10,888.8 (8682.8–13,471.6)	11,340.5	
Somalia	45.3 (8.4–127.5)	107.8 (24.8–251.0)	43.5 (9.5–112.7)	101.0 (24.2–242.5)	3058.6 (722.6–7430.6)	1761.52	
Sudan	96.5 (47.9–163.1)	158.6 (95.9–242.4)	101.0 (48.2–171.3)	146.7 (80.2–230.8)	21,814.0 (14,145.4–31,022.9)	10,515.2	
Syria	141.1 (88.4–207.5)	145.8 (99.7–205.0)	112.7 (67.3–164.2)	99.3 (67.9–131.0)	9987.0 (7397.8–12,880.3)	8109.9	
Tunisia	72.6 (37.9–111.7)	75.4 (49.4–105.8)	71.9 (37.2–118.8)	58.4 (34.7–87.3)	6256.6 (4163.4–8889.9)	8682.7	
United Arab Emirates	128.6 (76.3–188.5)	153.1 (100.1–214.7)	103.9 (60.2–156.2)	103.8 (68.8–148.4)	4460.4 (2941.6–6206.4)	1648.5	
Yemen	74.5 (27.2–149.7)	106.8 (43.0–205.8)	78.1 (31.1–158.7)	124.4 (59.2–225.4)	10,936.6 (5668.1–19,184.0)	5848.1	

Table 3 Disability-adjusted life years (DALYs) and years lived with disability (YLDs), with 95% uncertainty intervals (UI), per 100,000 population attributable to high body mass index among DALYs and YLDs due to ischemic heart disease, stroke, and diabetes, by age groups (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990 and 2015)

Age	Cause	DALYs						YLDs					
		1990			2015			1990			2015		
		Males	Females	Males	Females	Males	Females	Males	Females	Males	Females	Males	Females
15–49	Ischemic heart disease	473.0 (278.3–690.8)	328.0 (211.6–453.7)	560.1 (352.0–801.7)	327.5 (222.9–436.6)	5.9 (3.0–9.7)	5.3 (3.0–8.5)	9.9 (5.5–15.9)	8.9 (5.3–13.7)				
	Stroke	351.1 (225.4–483.7)	398.8 (286.0–519.7)	353.0 (234.5–484.9)	361.8 (261.7–468.2)	12.7 (7.0–20.1)	16.8 (10.5–24.9)	16.8 (9.9–25.6)	22.2 (14.2–31.4)				
	Diabetes	251.7 (163.0–351.2)	299.3 (216.5–399.8)	419.6 (284.2–577.4)	483.5 (351.3–640.4)	173.6 (103.8–262.9)	206.7 (133.2–297.0)	303.4 (187.6–448.4)	351.3 (229.4–491.8)				
50–69	Ischemic heart disease	3357.1 (1918.0–5002.9)	3270.6 (2220.9–4466.6)	3898.1 (2392.9–5519.0)	3014.3 (2122.1–3987.9)	74.0 (35.4–126.3)	73.6 (41.8–114.4)	116.6 (60.1–192.0)	105.3 (61.8–157.8)				
	Stroke	1686.2 (990.7–2460.1)	2247.9 (1556.8–3081.0)	1792.3 (1132.3–2514.2)	2015.4 (1440.7–2648.3)	64.4 (32.9–106.1)	87.3 (53.5–133.0)	81.1 (45.0–126.4)	99.5 (62.9–146.0)				
	Diabetes	1471.0 (909.2–2097.8)	2091.7 (1497.4–2815.7)	2417.9 (1604.3–3312.3)	3190.4 (2388.5–4121.2)	787.0 (431.0–1229.8)	1098.0 (689.5–1605.4)	1330.1 (769.0–1994.0)	1736.2 (1119.0–2482.3)				
70+	Ischemic heart disease	4001.4 (2028.0–6420.1)	4951.1 (3092.1–7158.9)	4600.8 (2513.9–7234.6)	4690.5 (2977.1–6737.5)	92.0 (42.0–167.8)	110.2 (61.8–180.6)	138.4 (66.9–240.6)	155.1 (88.7–248.2)				
	Stroke	1789.7 (919.0–2943.9)	2449.0 (1480.2–3678.2)	1875.5 (991.2–2972.6)	2200.4 (1362.7–3160.4)	60.1 (27.2–106.5)	78.8 (43.8–128.7)	72.2 (34.3–123.3)	90.8 (53.2–142.7)				
	Diabetes	1342.6 (725.2–2113.3)	1933.4 (1197.4–2848.9)	2345.0 (1336.8–3535.0)	3392.0 (2234.9–4736.3)	532.3 (258.2–920.5)	802.8 (468.0–1261.9)	921.4 (471.6–1515.4)	1307.7 (782.1–1998.0)				

Table 4 Disability-adjusted life years, with 95% uncertainty intervals (UI) per 100,000 population due to high body mass index observed in 1990 and 2015, and expected based on Socio-demographic Index in 2015 (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1990 and 2015)

Country	1990				2015				Total observed		Total expected
	Males		Females		Males		Females		Total observed	Total expected	
	Males	Females	Males	Females	Males	Females					
Afghanistan	4767.5 (2400.1–7893.8)	8437.9 (4934.2–12,974.6)	4677.5 (2411.1–7598.2)	8491.9 (5022.7–13,139.8)	965,879.5 (648,967.1–1,371,214.6)	206,389.0					
Bahrain	4039.9 (2769.8–5453.6)	4907.8 (3800.7–6091.1)	3109.6 (2127.8–4224.1)	3177.8 (2414.8–4028.7)	29,225.3 (22,628.0–36,438.7)	18,576.5					
Djibouti	1189.7 (362.1–2830.1)	1023.3 (378.0–2220.0)	2974.8 (1138.0–6633.1)	2633.4 (1192.2–5883.4)	15,630.0 (7793.1–29,440.4)	10,761.1					
Egypt	4317.3 (2880.3–5717.9)	4251.7 (3367.6–5153.0)	5025.6 (3688.3–6374.7)	4435.2 (3623.2–5284.9)	3,037,204.4 (2,478,932.1–3,605,791.4)	1,666,166.5					
Iran	2509.4 (1318.4–3910.9)	2733.1 (1877.6–3632.2)	2846.6 (1712.4–4306.3)	2696.7 (1880.0–3776.6)	1,698,192.1 (1,199,771.9–2,259,245.3)	1,493,421.0					
Iraq	6502.5 (4537.2–8970.9)	6404.8 (4660.0–8287.8)	6321.0 (4108.7–9096.4)	5726.9 (4046.8–7892.3)	1,090,796.9 (821,609.9–1,421,193.6)	452,426.9					
Jordan	4299.2 (2938.7–5943.2)	5362.9 (4166.0–6773.0)	3633.7 (2593.7–4668.3)	3176.4 (2469.7–3918.3)	141,869.1 (112,808.2–172,138.2)	106,862.3					
Kuwait	2861.0 (2115.1–3562.8)	3229.1 (2627.2–3868.5)	3026.5 (2239.0–3850.2)	2715.3 (2167.4–3389.3)	69,567.5 (55,117.3–84,812.6)	32,170.1					
Lebanon	3664.5 (2191.8–5471.8)	3454.1 (2306.5–4827.3)	2251.5 (1351.6–3307.2)	2480.0 (1658.6–3401.0)	123,560.0 (89,344.0–160,488.7)	110,981.5					
Libya	2379.4 (1551.1–3309.5)	3037.5 (2274.4–3882.7)	2904.3 (1926.1–3953.5)	3285.7 (2464.7–4171.7)	139,878.3 (108,615.2–173,319.3)	116,450.8					
Morocco	2490.8 (1420.6–3677.6)	2952.1 (1987.8–4068.0)	2336.9 (1332.7–3654.0)	3032.9 (1949.9–4325.8)	774,701.5 (555,295.0–1,031,236.8)	600,106.4					
Oman	2471.6 (1506.0–3723.1)	2805.3 (1835.8–4077.2)	2927.1 (1965.5–3972.0)	3097.2 (2308.7–3934.5)	75,807.5 (57,042.8–96,463.6)	64,799.1					
Pakistan	1543.7 (657.2–2752.2)	1869.4 (995.8–3001.1)	3009.9 (1619.5–4710.8)	3083.7 (1950.6–4398.8)	3,608,879.0 (2,420,204.5–4,972,204.0)	2,240,433.8					
Palestine	3235.1 (1896.7–4980.2)	3467.5 (2290.6–5022.7)	4300.8 (2694.9–6204.2)	3206.5 (2240.2–4368.5)	82,929.2 (61,735.5–109,791.4)	54,335.0					
Qatar	4074.6 (3099.9–5097.3)	5538.0 (4495.6–6656.9)	3236.7 (2297.7–4272.4)	3669.8 (2825.1–4589.9)	36,660.9 (27,986.6–46,001.8)	23,177.6					
Saudi Arabia	2296.3 (1512.6–3138.8)	2558.8 (1946.2–3250.2)	2507.0 (1808.4–3262.0)	2165.4 (1693.7–2699.8)	490,128.6 (383,013.6–601,300.7)	440,497.5					
Somalia	1463.9 (330.5–3897.1)	2980.2 (780.2–6721.4)	1325.4 (330.3–3416.1)	2828.7 (834.7–6943.8)	102,092.6 (31,197.6–246,633.4)	59,469.2					
Sudan	2883.9 (1513.0–4693.9)	4378.6 (2790.2–6417.8)	3057.9 (1567.6–4893.5)	4128.4 (2480.3–6203.9)	764,380.1 (523,579.9–1,052,574.8)	364,535.7					
Syria	4198.1 (2692.6–5980.2)	4090.6 (2917.9–5509.0)	3305.8 (2135.7–4629.7)	2889.2 (2118.5–3677.0)	347,354.1 (267,569.5–437,633.2)	276,598.4					
Tunisia	2065.4 (1134.9–3084.8)	2248.3 (1594.2–2947.4)	2165.4 (1242.3–3352.4)	1889.1 (1269.8–2570.8)	217,846.2 (151,390.5–292,471.8)	272,878.3					
United Arab Emirates	4028.4 (2596.4–5648.6)	4675.9 (3270.4–6389.7)	3425.1 (2209.4–4864.9)	3491.9 (2522.1–4661.1)	242,427.0 (171,328.5–322,519.2)	87,802.7					
Yemen	2247.0 (881.5–4390.2)	3017.4 (1304.3–5529.8)	2373.0 (1016.2–4579.6)	3639.9 (1893.9–6356.4)	393,538.0 (216,892.2–664,897.8)	207,048.3					

measured by SDI (Ulijaszek 2007). Cultural factors and obesogenic cultural traditions disproportionately affect women, highlighting cross-cutting gender issues in women's health (Shapira 2013). This might explain the observed gender differences in overweight and obesity. Similarly, social determinants of health in countries in conflict tend to differ as morbidity and mortality are associated with conflict (World Health Organization 2008). More importantly, wars and civil unrest are closely associated with child malnutrition, which in turn is associated with increased risk of obesity, hypertension, cardiovascular disease, and type 2 diabetes (Devakumar et al. 2014; Charchuk et al. 2015). This leads to the intergenerational effects of war on obesity and the doubled burden of undernutrition in countries affected by wars or experiencing chronic civil unrest (Devakumar et al. 2014).

Despite this increase in obesity, EMR countries can hope to control their epidemic as declines in obesity prevalence have been reported in other countries due to health interventions and environmental and policy changes (Schmidt Morgen et al. 2013; Keane et al. 2014). EMR countries can benefit from implementing an array of proven interventions to control their obesity epidemics. Over the last decade, very few EMR ministries of health have focused on health promotion strategies to reduce obesity, such as awareness and behavioral changes. Of the 22 countries in the EMR, only Bahrain, Qatar, and Saudi Arabia had substantial information on combating obesity on their ministry of health websites (Gharib et al. 2012; Saudi Ministry of Health 2012). Qatar and Saudi Arabia launched national campaigns against obesity in 2011 and 2012, respectively. The Saudi campaign focused primarily on dietary awareness, promoting healthy eating choices and emphasizing variety and balance through a food pyramid (Hamad Al-Dkheel 2012). The Qatar campaign was more comprehensive, setting up programs to increase physical activity, replace school snacks with healthier options, and pilot a nutrition surveillance system in addition to a dietary awareness campaign (Mohammed Al-Thani 2011). Indeed, and despite the campaign in Qatar, obesity prevalence is one of the highest in the region, suggesting the need for a more aggressive approach.

Promoting healthier lifestyles around nutrition and physical activity is greatly needed in this region. As pointed out previously, possible factors determining obesity in the EMR include nutrition transition, inactivity, urbanization, marital status, a shorter duration of breastfeeding, frequent snacking, skipping breakfast, a high intake of sugary beverages, an increase in the incidence of eating outside the home, long periods of time spent viewing television, massive marketing promotion of high-fat foods, stunting, perceived body image, cultural elements, and food subsidy policies (Musaiger 2011b).

Indeed, estimates from the GBD study show an increase in dietary risk factors and low levels of physical activity (Forouzanfar et al. 2016). What the EMR populations consume can be directly affected by national policies, specifically those around food economics. For instance, in many EMR countries such as Egypt, Morocco, and Saudi Arabia, governments subsidize grains, bread, wheat flour, sugar, and cooking oil, and impose taxes on imported foods (Asfaw 2007; Musaiger 2011b). However, fruits and vegetables are neither subsidized nor exempt from import taxes, making healthier food choices harder for the population. As for physical activity, the lack of exercise facilities has been reported, at least in Egypt and Saudi Arabia, as the main reason for low physical activity (Al-Rafaae and Al-Hazzaa 2001; El-Gilany et al. 2011).

This study has a few limitations. These include possible underestimation of the prevalence of obesity or disease burden from high BMI as self-reported height and weight data were used. However, we corrected these based on measured data at each age, sex, and country unit (GBD 2015 Obesity Collaborators 2017). Briefly, no significant difference between measured and self-reported data for children was found. For adults, self-reported data were adjusted for overweight prevalence, obesity prevalence, and mean BMI using nested hierarchical mixed-effects regression models, fit using restricted maximum likelihood separately by sex. Second, estimates for some countries with sparse data were driven by covariates in the statistical modeling. This, in fact, highlights the need for more timely surveillance data on BMI and risk factors in the EMR. Better quality data in the EMR for quantification of morbidity and mortality burden and health policy planning is urgently needed. Third, the attributable effect of BMI on ischemic heart disease, stroke, and diabetes was derived from prospective observational studies and meta-analyses, and hence does not account for differences by ethnicity, or for underlying diseases (Global BMI Mortality Collaboration et al. 2016). While these observational studies are not specifically from the EMR, the attributable effect of BMI on the three diseases has been shown to remain the same across different regions from the world (Singh et al. 2013). Fourth, BMI presents some drawbacks as it is a measure of excess weight rather than excess body fat (Daniels 2009). The relationship between BMI and body fat can be influenced by age, sex, ethnicity, and muscle mass. In addition, BMI does not indicate whether excess weight is due to fat, muscle, or bone (Daniels 2009). However, BMI is favored for its ease of use in surveys and is the most available indicator regarding excess weight. More details on this study's limitations are available elsewhere (GBD 2015 Obesity Collaborators 2017).

This study showed that high BMI creates a major burden in the EMR. We call for countries in the region to invest

more resources in prevention and health promotion efforts to reduce the prevalence of obesity. These programs should focus on stopping weight gain as a first step and more aggressive programs to reduce weight among those who need to do so. These programs should take into account the culture and local environment. Moreover, countries should join efforts in their programs and policies and share experiences and success stories.

GBD 2015 Eastern Mediterranean Region Obesity Collaborators:

Ali H. Mokdad, PhD (corresponding author), Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Charbel El Bcheraoui, PhD, Institute for Health Metrics and Evaluation, University of Washington. Ashkan Afshin, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, United States. Raghid Charara, MD, American University of Beirut, Beirut, Lebanon. Ibrahim Khalil, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Maziar Moradi-Lakeh, MD, Department of Community Medicine, Preventive Medicine and Public Health Research Center, Gastrointestinal and Liver Disease Research Center (GILDRC), University of Medical Sciences, Tehran, Iran. Nicholas J. Kassebaum, MD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States; Department of Anesthesiology & Pain Medicine, Seattle Children's Hospital, Seattle, Washington, United States. Michael Collison, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Farah Daoud, BA/BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Kristopher J. Krohn, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Adrienne Chew, ND, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Stan H. Biryukov, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Leslie Cornaby, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Kyle J. Foreman, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States; Imperial College London, London, United Kingdom. Michael Kutz, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Patrick Liu, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Marissa Reitsma, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Patrick Sur, BA, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States, Seattle. Haidong Wang, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Ben Zipkin, BS, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States, Seattle. Johan Ärnlöv, PhD, Department of Neurobiology, Care Sciences and Society, Division of Family Medicine and Primary Care, Karolinska Institutet, Stockholm, Sweden; School of Health and Social Studies, Dalarna University, Falun, Sweden. Cristiana Abbafati, PhD, La Sapienza, University of Rome, Rome, Italy. Abdishakur M. Abdulle, PhD, New York University Abu Dhabi, Abu Dhabi, United Arab Emirates. Niveen M.E. Abu-Rmeileh, PhD, Institute of Community and Public Health, Birzeit University, Ramallah, West Bank, Palestine. Muktar Beshir Ahmed, MPH, College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator, Jimma University, Jimma, Oromiya, Ethiopia. Ziyad Al-Aly, MD, Washington University in St. Louis, St. Louis, MO, United States. Khurshid Alam, PhD,

Murdoch Childrens Research Institute, The University of Melbourne, Parkville, Victoria, Australia; The University of Melbourne, Melbourne, VIC, Australia; The University of Sydney, Sydney, NSW, Australia. Reza Alizadeh-Navaei, PhD, Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Mazandaran, Iran. Ala'a Alkerwi, PhD, Luxembourg Institute of Health (LIH), Strassen, Luxembourg. Khalid A. Altirkawi, MD, King Saud University, Riyadh, Saudi Arabia. Nelson Alvis-Guzman, PhD, Universidad de Cartagena, Cartagena de Indias, Colombia. Bernhard T. Baune, PhD, School of Medicine, University of Adelaide, Adelaide, South Australia, Australia. Neeraj Bedi, MD, College of Public Health and Tropical Medicine, Jazan, Saudi Arabia. Derrick A. Bennett, PhD, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom. Addisu S. Beyene, MPH, College of Health and Medical Science, Haramaya University, Harar, Ethiopia. Zulfiqar A. Bhutta, PhD, Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan; Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON, Canada. Mulugeta M. Birhanu, MS, University of Groningen, UMCG, Groningen, Groningen, Netherlands; Mekelle University, Mekelle, Ethiopia. Hadi Danawi, PhD, Walden University, Minneapolis, Minnesota, United States. Seyed-Mohammad Fereshtehnejad, PhD, Department of Neurobiology, Care Sciences and Society (NVS), Karolinska Institutet, Stockholm, Sweden. Florian Fischer, PhD, School of Public Health, Bielefeld University, Bielefeld, Germany. Tsegaye Tewelde Gebrehiwot, MPH, Jimma University, Jimma, Oromia, Ethiopia. Paramjit Singh Gill, DM, Warwick Medical School, University of Warwick, Coventry, United Kingdom; University of Birmingham, UK, Birmingham, United Kingdom. Philimon N. Gona, PhD, University of Massachusetts Boston, Boston, Massachusetts, United States. Vipin Gupta, PhD, Department of Anthropology, University of Delhi, Delhi, Delhi, India. Tesfa Dejenie Habtewold, MS, University of Groningen, Groningen, Netherlands; Debre Berhan University, Debre Berhan, Ethiopia. Randah Ribhi Hamadeh, DPhil, Arabian Gulf University, Manama, Bahrain. Samer Hamidi, DrPH, Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates. Habtamu Abera Hareri, MS, Addis Ababa University, Addis Ababa, Ethiopia. Masako Horino, MPH, Bureau of Child, Family & Community Wellness, Nevada Division of Public and Behavioral Health, Carson City, NV, United States. Mohamed Hsairi, MD, Department of Epidemiology, Salah Azaiz Institute, Tunis, Tunisia. Mehdi Javanbakht, PhD, University of Aberdeen, Aberdeen, Aberdeen, United Kingdom. Denny John, MPH, International Center for Research on Women, New Delhi, Delhi, India. Jost B. Jonas, MD, Department of Ophthalmology, Medical Faculty Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany. Vasna Joshua, PhD, National Institute of Epidemiology, Indian Council of Medical Research, Chennai, India. Amir Kasaeian, PhD, Hematology-Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran; Endocrinology and Metabolism Population Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran. Ezra B. Ketema, MS, Mekelle University, Mekelle, Ethiopia. Yousef Saleh Khader, ScD, Department of Community Medicine, Public Health and Family Medicine, Jordan University of Science and Technology, Irbid, Irbid, Jordan. Ejaz Ahmad Khan, MD, Health Services Academy, Islamabad, Punjab, Pakistan. Jagdish Khubchandani, PhD, Department of Nutrition and Health Science, Ball State University, Muncie, Indiana, United States. Daniel Kim, DrPH, Department of Health Sciences, Northeastern University, Boston, Massachusetts, United States. Yun Jin Kim, PhD, Faculty of Chinese Medicine, Southern University College, Skudai, Johor, Malaysia. Yohannes Kinfu, PhD, Centre for Research and Action in Public Health, University of Canberra, Canberra, Australian Capital Territory, Australia. Yoshihiro Kokubo, PhD, Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Osaka,

Japan. Heidi J. Larson, PhD, Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Paul H. Lee, PhD, Hong Kong Polytechnic University, Hong Kong, China. Raimundas Lunevicius, PhD, Aintree University Hospital National Health Service Foundation Trust, Liverpool, United Kingdom; School of Medicine, University of Liverpool, Liverpool, United Kingdom. Hassan Magdy Abd El Razek, MBChB, Mansoura Faculty of Medicine, Mansoura, Egypt. Mohammed Magdy Abd El Razek, MBChB, Aswan University Hospital, Aswan Faculty of Medicine, Aswan, Egypt. Reza Malekzadeh, MD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran. Kedar K. Mate, MSc, McGill University, Montreal, Quebec, Canada. Ziad A. Memish, MD, Saudi Ministry of Health, Riyadh, Saudi Arabia; College of Medicine, Alfaisal University, Riyadh, Saudi Arabia. Mubarek Abera Mengistie, MS, Jimma University, Jimma, Oromia, Ethiopia. George A. Mensah, MD, Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, United States. Felix Akpojone Ogbo, MPH, Centre for Health Research, Western Sydney University, Sydney, New South Wales, Australia. Farshad Pourmalek, PhD, University of British Columbia, Vancouver, British Columbia, Canada. Anwar Rafay, MS, Contech International Health Consultants, Lahore, Punjab, Pakistan; Contech School of Public Health, Lahore, Punjab, Pakistan. Mostafa Qorbani, PhD, Non-communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran. Vafa Rahimi-Movaghar, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Tehran, Iran. Saleem M. Rana, PhD, Contech School of Public Health, Lahore, Punjab, Pakistan; Contech International Health Consultants, Lahore, Punjab, Pakistan. Salman Rawaf, MD, Imperial College London, London, United Kingdom. Andre M.N. Renzaho, PhD, Western Sydney University, Penrith, NSW, NSW, Australia. Satar Rezaei, PhD, School of Public Health, Kermanshah University of Medical Sciences, Kermanshah, Iran. Mohammad Sadegh Rezaei, MD, Infectious Disease Research Centre with Focus on Nosocomial Infection, Mazandaran University of Medical Sciences, Sari, Mazandaran, Iran. Mahdi Safdarian, MD, Sina Trauma & Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran. Mohammad Ali Sahraian, MD, MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran. Payman Salamati, MD, Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Tehran, Iran. Abdallah M. Samy, PhD, Ain Shams University, Cairo, Egypt. Juan Ramon Sanabria, MD, J Edwards School of Medicine, Marshall University, Huntington, WV, United States; Case Western Reserve University, Cleveland, OH, United States. Milena M. Santric Milicevic, PhD, Institute of Social Medicine, Faculty of Medicine, University of Belgrade, Serbia, Serbia; Centre School of Public Health and Health Management, Faculty of Medicine, University of Belgrade, Serbia, Serbia. Benn Sartorius, PhD, Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa; UKZN Gastrointestinal Cancer Research Centre, South African Medical Research Council (SAMRC), Durban, South Africa. Sadaf G. Sepanlou, PhD, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Tehran, Iran. Masood Ali Shaikh, MD, Independent Consultant, Karachi, Pakistan. Rahman Shiri, PhD, Finnish Institute of Occupational Health, Work Organizations, Work Disability Program, Department of Public Health, Faculty of Medicine, University of Helsinki, Helsinki, Finland. Diego Augusto Santos Silva, PhD, Federal University of Santa Catarina, Florianopolis, Brazil. Jasvinder A. Singh, MD, University of Alabama at Birmingham and Birmingham Veterans Affairs Medical Center, Birmingham, Alabama, United States. Abdullah Sulieman Terkawi, MD, Department of

Anesthesiology, University of Virginia, Charlottesville, VA, United States; Department of Anesthesiology, King Fahad Medical City, Riyadh, Saudi Arabia; Outcomes Research Consortium, Cleveland Clinic, Cleveland, OH, United States. Tenaw Yimer Tiruye, MPH, Debre Markos University, Debre Markos, Ethiopia. Kingsley Nnanna Ukwaja, MD, Department of Internal Medicine, Federal Teaching Hospital, Abakaliki, Ebonyi State, Nigeria. Tommi Vasankari, PhD, UKK Institute for Health Promotion Research, Tampere, Finland. Vasily Victorovich Vlassov, MD, National Research University Higher School of Economics, Moscow, Russia. Tolassa Wakayo, MS, Jimma University, Jimma, Oromia, Ethiopia. Andrea Werdecker, PhD, Competence Center Mortality-Follow-Up of the German National Cohort, Federal Institute for Population Research, Wiesbaden, Hesse, Germany. Naohiro Yonemoto, MPH, Department of Biostatistics, School of Public Health, Kyoto University, Kyoto, Japan. Mustafa Z. Younis, DrPH, Jackson State University, Jackson, MS, United States. Bassel Zein, MS, Department of Neuroscience, Georgetown University, Washington DC, United States. Aisha O. Jumaan, PhD, Independent Consultant, Seattle, Washington, United States. Theo Vos, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Simon I. Hay, DSc, Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, United Kingdom; Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Mohsen Naghavi, PhD, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States. Christopher J. L. Murray, DPhil, Institute for Health Metrics and Evaluation, University of Washington, Seattle, Washington, United States.

Compliance with ethical standards

This manuscript reflects original work that has not previously been published in whole or in part and is not under consideration elsewhere. All authors have read the manuscript and have agreed that the work is ready for submission and accept responsibility for its contents. The authors of this paper have complied with all ethical standards and do not have any conflicts of interest to disclose at the time of submission. The funding source played no role in the design of the study, the analysis and interpretation of data, and the writing of the paper. The study did not involve human participants and/or animals; therefore, no informed consent was needed.

Funding This research was funded by the Bill & Melinda Gates Foundation.

Conflict of interest The authors declare that they have no conflicts of interest at this time.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abou Abbas L, Salameh P, Nasser W et al (2015) Obesity and symptoms of depression among adults in selected countries of the Middle East: a systematic review and meta-analysis. *Clin Obes* 5:2–11. doi:10.1111/cob.12082

- Al-Rafae SA, Al-Hazzaa HM (2001) Physical activity profile of adult males in Riyadh city. *Saudi Med J* 22:784–789
- Asfaw A (2007) Do government food price policies affect the prevalence of obesity? Empirical evidence from Egypt. *World Dev* 35:687–701. doi:10.1016/j.worlddev.2006.05.005
- Broyles ST, Denstel KD, Church TS et al (2015) The epidemiological transition and the global childhood obesity epidemic. *Int J Obes Suppl* 5:S3–S8. doi:10.1038/ijosup.2015.12
- Charchuk R, Houston S, Hawkes MT (2015) Elevated prevalence of malnutrition and malaria among school-aged children and adolescents in war-ravaged South Sudan. *Pathog Glob Health* 109:395–400. doi:10.1080/20477724.2015.1126033
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 320:1240
- Daniels SR (2009) The use of BMI in the clinical setting. *Pediatrics* 124(Suppl 1):S35–S41. doi:10.1542/peds.2008-3586F
- Devakumar D, Birch M, Osrin D et al (2014) The intergenerational effects of war on the health of children. *BMC Med* 12:57. doi:10.1186/1741-7015-12-57
- Drewnowski A, Moudon AV, Jiao J et al (2014) Food environment and socioeconomic status influence obesity rates in Seattle and in Paris. *Int J Obes* 38:306–314. doi:10.1038/ijo.2013.97
- El-Gilany AH, Badawi K, El-Khawaga G, Awadalla N (2011) Physical activity profile of students in Mansoura University, Egypt. *East Mediterr Health J Rev Sante Mediterr Orient Al-Majallah Al-Sihhiyah Li-Sharq Al-Mutawassit* 17:694–702
- Forouzanfar MH, Alexander L, Anderson HR et al (2015) Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 386:2287–2323. doi:10.1016/S0140-6736(15)00128-2
- Forouzanfar MH, Afshin A, Alexander LT et al (2016) Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1659–1724. doi:10.1016/S0140-6736(16)31679-8
- GBD 2015 Obesity Collaborators (2017) Health effects of overweight and obesity in 195 Countries over 25 Years. *N Engl J Med* 377:13–27. doi:10.1056/NEJMoa1614362
- Gharib N, Al Amer M, Al-Salehi S (2012) Nutrition clinics: management and prevention of obesity. http://hajj.moh.gov.bh/pdf/publications/X_217201413620.pdf. Accessed 27 Apr 2017
- Global BMI Mortality Collaboration null, Di Angelantonio E, Bhupathiraju S et al (2016) Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet Lond Engl* 388:776–786. doi:10.1016/S0140-6736(16)30175-1
- Hamad Al-Dkheel M (2012) Dietary Guidelines for Saudis: the healthy food palm. <http://www.moh.gov.sa/en/Ministry/MediaCenter/Publications/Documents/final%20english%20%20%D8%A7%D9%84%D9%83%D8%AA%D8%A7%D8%A8%20%D8%A7%D9%84%D8%B9%D9%84%D9%85%D9%8A%20%D8%A5%D9%86%D8%AC%D9%84%D9%8A%D8%B2%D9%8A.pdf>. Accessed 27 Apr 2017
- Hill AB (1965) The environment and disease: association or causation? *Proc R Soc Med* 58:295–300
- Institute for Health Metrics and Evaluation (2016) Global Burden of Disease Study 2015 (GBD 2015) Data Resources | GHDx. <http://ghdx.healthdata.org/gbd-2015>. Accessed 30 May 2017
- Keane E, Kearney PM, Perry IJ et al (2014) Trends and prevalence of overweight and obesity in primary school aged children in the Republic of Ireland from 2002–2012: a systematic review. *BMC Public Health* 14:974. doi:10.1186/1471-2458-14-974
- Kelly SP, Graubard BI, Andreotti G et al (2017) Prediagnostic body mass index trajectories in relation to prostate cancer incidence and mortality in the PLCO cancer screening trial. *J Natl Cancer Inst* 109:1–9. doi:10.1093/jnci/djw225
- Kim SH, Després J-P, Koh KK (2016) Obesity and cardiovascular disease: friend or foe? *Eur Heart J* 37:3560–3568. doi:10.1093/eurheartj/ehv509
- Maciosek MV, LaFrance AB, Dehmer SP et al (2017) Updated priorities among effective clinical preventive services. *Ann Fam Med* 15:14–22. doi:10.1370/afm.2017
- Mehta NK, Patel SA, Ali MK, Venkat Narayan KM (2017) Preventing disability: the influence of modifiable risk factors on state and national disability prevalence. *Health Aff Proj Hope* 36:626–635. doi:10.1377/hlthaff.2016.1281
- Memish ZA (2014) Obesity and associated factors—Kingdom of Saudi Arabia, 2013. *Prev Chronic Dis*. doi:10.5888/pcd11.140236
- Mohammed Al-Thani AA (2011) Qatar national nutrition and physical activity action plan 2011–2016. <https://extranet.who.int/nutrition/gina/sites/default/files/QAT%202011%20National%20Nutrition%20and%20Physical%20Activity%20Action%20Plan.pdf>. Accessed 27 Apr 2017
- Mokdad AH, Forouzanfar MH, Daoud F et al (2016) Health in times of uncertainty in the eastern Mediterranean region, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Glob Health* 4:e704–e713. doi:10.1016/S2214-109X(16)30168-1
- Musaiger AO (2011a) Food consumption patterns in the Eastern Mediterranean Region. https://www.researchgate.net/publication/236668592_Food_Consumption_Patterns_in_the_Eastern_Mediterranean_Region. Accessed 27 Apr 2017
- Musaiger AO (2011b) Overweight and obesity in eastern mediterranean region: prevalence and possible causes. *J Obes* 2011:407237. doi:10.1155/2011/407237
- Musaiger AO, Al-Hazzaa HM, Takruri HR, Mokhtar N (2012) Change in nutrition and lifestyle in the Eastern Mediterranean Region: health Impact. *J Nutr Metab* 2012:e436762. doi:10.1155/2012/436762
- Ng M, Fleming T, Robinson M et al (2014) Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 384:766–781. doi:10.1016/S0140-6736(14)60460-8
- Ng M, Liu P, Thomson B, Murray CJL (2016) A novel method for estimating distributions of body mass index. *Popul Health Metr* 14:6. doi:10.1186/s12963-016-0076-2
- Pratt LA, Brody DJ (2014) Depression and obesity in the U.S. adult household population, 2005–2010. *NCHS Data Brief* 167:1–8
- Saudi Ministry of Health (2012) National campaign against overweight and obesity. <http://www.moh.gov.sa/en/>. Accessed 7 Apr 2017
- Schmidt Morgen C, Rokholm B, Sjöberg Brixval C et al (2013) Trends in prevalence of overweight and obesity in danish infants, children and adolescents—are we still on a plateau? *PLoS One* 8:e69860. doi:10.1371/journal.pone.0069860
- Shapira N (2013) Women’s higher health risks in the obesogenic environment: a gender nutrition approach to metabolic dimorphism with predictive, preventive, and personalised medicine. *EPMA J* 4:1. doi:10.1186/1878-5085-4-1
- Sibai AM, Hwalla N, Adra N, Rahal B (2003) Prevalence and covariates of obesity in lebanon: findings from the first epidemiological study. *Obes Res* 11:1353–1361. doi:10.1038/oby.2003.183

- Sidney S, Quesenberry CP, Jaffe MG et al (2016) Recent trends in cardiovascular mortality in the United States and public health goals. *JAMA Cardiol* 1:594–599. doi:[10.1001/jamacardio.2016.1326](https://doi.org/10.1001/jamacardio.2016.1326)
- Singh GM, Danaei G, Farzadfar F et al (2013) The age-specific quantitative effects of metabolic risk factors on cardiovascular diseases and diabetes: a pooled analysis. *PLoS One* 8:e65174. doi:[10.1371/journal.pone.0065174](https://doi.org/10.1371/journal.pone.0065174)
- Stevens GA, Alkema L, Black RE et al (2016) Guidelines for accurate and transparent health estimates reporting: the GATHER statement. *Lancet* 388:e19–e23. doi:[10.1016/S0140-6736\(16\)30388-9](https://doi.org/10.1016/S0140-6736(16)30388-9)
- Terkawi AS, Rafiq M, Algadaan R et al (2014) General anesthesia for the heaviest man in the world. *Saudi J Anaesth* 8:S101–S104. doi:[10.4103/1658-354X.144087](https://doi.org/10.4103/1658-354X.144087)
- Ulijaszek SJ (2007) Frameworks of population obesity and the use of cultural consensus modeling in the study of environments contributing to obesity. *Econ Hum Biol* 5:443–457. doi:[10.1016/j.ehb.2007.08.006](https://doi.org/10.1016/j.ehb.2007.08.006)
- Wang H, Naghavi M, Allen C et al (2016) Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388:1459–1544. doi:[10.1016/S0140-6736\(16\)31012-1](https://doi.org/10.1016/S0140-6736(16)31012-1)
- World Cancer Research Fund, American Institute for Cancer Research (2007) Food, nutrition, physical activity, and the prevention of cancer: a global perspective. AICR, Washington, DC
- World Health Organization (2008) Integrated health services—what and why? http://www.who.int/healthsystems/technical_brief_final.pdf. Accessed 27 Apr 2017
- World Health Organization (2016) Eastern Mediterranean Region office. About us. <http://www.emro.who.int/entity/about-us/index.html>. Accessed 27 Apr 2017
- Yao T-C, Tsai H-J, Chang S-W et al (2017) Obesity disproportionately impacts lung volumes, airflow and exhaled nitric oxide in children. *PLoS One* 12:e0174691. doi:[10.1371/journal.pone.0174691](https://doi.org/10.1371/journal.pone.0174691)

Title: Burden of obesity in the Eastern Mediterranean Region: findings from the Global Burden of Disease 2015 study

Group name: GBD 2015 Eastern Mediterranean Region Obesity Collaborators

Prefix	First Name	Middle Name	Last Name	Address	Highest Degree	Department	Institution/Organization	Institution/Organization City	Institution/Organization State	Institution/Organization 2	Institution/Organization 2 City	Institution/Organization 2 State	Institution/Organization 2	Institution	Institution	Institution	
Dr.	Charbel		El Bcharao	charbel@uw.edu		Institute for Health Metrics and Evaluation	University of Washington										
Dr.	Ashkan		Afshin	ashkan.afsi@MD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	WA			United_States					
Dr.	Raghd		Charara	raghdrc@gm			American University of Beirut	Beirut				Lebanon					
Dr.	Ibrahim		Khalil	ikhallil@uw.edu		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Dr.	Maziar		Moradi-Lal	lamazmorad MD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	WA			United_States	Department of Iran University of Medical Sciences	Tehran			Iran
Dr.	Nicholas	J	Kassebaurn	nickjk@uw MD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States	Department of Seattle Children's Hospital	Seattle			United_States
Mr.	Michael		Collison	mic314@uw.edu		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Ms.	Farah		Daoud	farahd@uw.edu		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Kristopher	J.	Krohn	kkrohn1@BA		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Ms.	Adrienne		Chew	apchew@ND		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Stan	H.	Biryukov	stan0625@BS		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Ms.	Leslie		Cornaby	lcomnaby@BS		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Kyle	J.	Foreman	kfor@uw.e PhD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States	Imperial College London	London			United_Kingdom
Mr.	Michael		Kutz	kutz13@uw BS		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Patrick		Liu	pyliu@uw BA		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Ms.	Marissa		Reitsma	mreitsma@BS		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Patrick		Sur	psur2417@BA		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States		Seattle			
Dr.	Haidong		Wang	haidong@PhD		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States					
Mr.	Ben		Zipkin	zipkinb@uw.edu		Institute for Health Metrics and Evaluation	University of Washington	Seattle	Washington			United_States		Seattle			
						Department of Neurobiology, Care Sciences and Society, Division of Family Medicine and Primary Care											
Prof.	Johan		Ärnölv	johan.arnic@PhD			Karolinska Institutet	Stockholm				Sweden	School of Uppsala University	Falun			Sweden
Dr.	Cristiana		Abbafati	cristiana.al@PhD			La Sapienza, University of Rome	Rome				Italy					
Dr.	Abdshakur	M	Abdulle	aa192@ny PhD			New York University Abu Dhabi	Abu Dhabi				United Arab Emirates					
Dr.	Niveen	M.E.	Abu-Rmeili	nrmelleh@PhD		Institute of Community and Public Health	Birzeit University	Ramallah				West Bank	Palestine				
						College of Health Sciences, Department of Epidemiology, ICT and e-Learning Coordinator											
Mr.	Muktar	Beshir	Ahmed	muktar27@ MPH			Jimma University	Jimma				Oromiya					
Dr.	Ziyad		Al-Aly	zalah1@gm MD			Washington University in St. Louis	St. Louis	MO			United_States					
Dr.	Khurshid		Alam	khurshid.b@PhD		Murdoch Childrens Research Institute	The University of Melbourne	Parville	Victoria			Australia	The University of Melbourne	Melbourne	VIC		Australia
Dr.	Reza		Alizadeh-N	reza_nava@PhD		Gastrointestinal Cancer Research Center	Mazandaran University of Medical Sciences	Sari	Mazandaran			Iran					
Dr.	Ala'a		Alkerwi	alaa.alkerw@PhD			Luxembourg Institute of Health (LIH)	Strassen				Luxembourg					
Dr.	Khalid	A.	Altirkawi	kaltirkawi@MD			King Saud University	Riyadh				Saudi_Arabia					
Prof.	Nelson		Alvis-Guzm	nalvis@yal PhD			Universidad de Cartagena	Cartagena	de Indias			Colombia					
Prof.	Bernhard	T	Baune	bernhard.t@PhD		School of Medicine	University of Adelaide	Adelaide				South Australia					Australia
Dr.	Neeraj		Bedi	drnbedi@yMD			College of Public Health and Tropical Medicine	Jazan				Saudi_Arabia					
Dr.	Derrick	A.	Bennett	derrick.ber@PhD		Nuffield Department of Population Health	University of Oxford	Oxford				United_Kingdom					
Mr.	Addisu	S.	Beyene		MPH	College of Health and Medical Science	Haramaya University	Harar				Ethiopia					
Dr.	Zulfiqar	A.	Bhutta	zulfiqar.bh@PhD		Centre of Excellence in Women and Child Health	Aga Khan University	Karachi				Pakistan	Centre for The Hospital for Sick Children	Toronto			Canada
Mr.	Mulugeta	M.	Birhanu	muler.wars@MS			University of Groningen, UMGCG	Groningen				Netherlands	Mekelle University	Mekelle			Ethiopia
Dr.	Hadi		Danawi	hdanawi@PhD			Walden University	Minneapolis				United_States					
						Department of Neurobiology, Care Sciences and Society (NVS)											
Dr.	Seyed-Mohammad		Fereshtehr	sm.feresh@PhD		School of Public Health	Karolinska Institutet	Stockholm				Sweden					
Dr.	Florian		Fischer		PhD		Bielefeld University	Bielefeld				Germany					
Mr.	Tsegaye	Tewelde	Gebrehiwo	tsegaye.ten@ MPH			Jimma University	Jimma				Oromia					
Prof.	Paramjit	Singh	Gilli	P.Gilli1@w@FRCGP		Warwick Medical School	University of Warwick	Coventry				United_Kingdom	University of Birmingham, UK	Birmingham			United_Kingdom
Prof.	Phillimon	N	Gona	phillimon@PhD			University of Massachusetts Boston	Boston				Massachusetts					
Dr.	Vipin		Gupta	udailig@grr@PhD		Department of Anthropology	University of Delhi	Delhi				India					
Mr.	Tesfa	Dejenie	Habtewold	tesfades@PhD			University of Groningen	Groningen				Netherlands	Debre Berhan University	Debre Berhan			Ethiopia
Prof.	Randah	Ribhi	Hamadeh	randah@aj@DPH			Arabian Gulf University	Manama				Bahrain					
Dr.	Samer		Hamidi	s.hamidi@DrPH			Hamdan Bin Mohammed Smart University	Dubai				United_Arab_Emirates					
Mr.	Habtamu	Abera	Hareri	habtamu.a@MS			Addis Ababa university	Addis Ababa				Ethiopia					
Ms.	Masako		Horino	mhorino@ MPH		Bureau of Child, Family & Community Wellness	Nevada Division of Public and Behavioral Health	Carson City	NV			United_States					
Prof.	Mohamed		Hsairi	mohamed.MD		Department of Epidemiology	Salah Azaiz Institute	Tunis				Tunisia					
Dr.	Mehdi		Javanbakt	javanbakt@PhD			University of Aberdeen	Aberdeen				United_Kingdom					
Mr.	Denny		John	djohn1976 MPH			International Center for Research on Women	New Delhi				India					
						Department of Ophthalmology, Medical Faculty Mannheim											
Prof.	Jost	B	Jonas	Jost.Jonas@MD			Ruprecht-Karls-University Heidelberg	Mannheim				Germany					
Dr.	Vasna		Joshua	vasna.joshi@PhD		National Institute of Epidemiology	Indian Council of Medical Research	Chennai				India					
						Hematology-Oncology and Stem Cell Transplantation Research Center											
Prof.	Amir		Kasaean		PhD		Tehran University of Medical Sciences	Tehran				Iran					
Mr.	Ezra	B.	Ketema		MS		Mekelle University	Mekelle				Ethiopia					Iran
						Department of Community Medicine, Public Health and Family Medicine											
Prof.	Yousef	Saleh	Khader	yskhader@ScD			Jordan University of Science and Technology	Irbid				Jordan					
Dr.	Ejaz	Ahmad	Khan	ejaz@hsa.i MD			Health Services Academy	Islamabad				Punjab					Pakistan
Dr.	Jagdish		Khubchanc	khubchanc@PhD		Department of Nutrition and Health Science	Ball State University	Muncie				Indiana					United_States
Dr.	Daniel		Kim			Department of Health Sciences,	Northeastern University	Boston				Massachusetts					United_States
Dr.	Yun Jin		Kim	neurokim7@PhD		Faculty of Chinese Medicine	Southern University College	Skudai				Johor					Malaysia
Dr.	Yohannes		Kinfu	yohannes.i@PhD		Centre for Research and Action in Public Health	University of Canberra	Canberra				Australian Capital Territory					Australia
Dr.	Yoshihiro		Kokubo	ykokubo@PhD		Department of Preventive Cardiology	National Cerebral and Cardiovascular Center	Suita				Osaka					Japan
						Department of Infectious Disease Epidemiology											
Dr.	Heidi	J.	Larson	Heidi.Larsc@PhD			London School of Hygiene & Tropical Medicine	London				United_Kingdom	Institute for University of Washington	Seattle			United_States
Dr.	Paul	H	Lee	paul.h.lee@PhD			Hong Kong Polytechnic University	Hong Kong				China					
							Aintree University Hospital National Health Service										
Prof.	Raimundas		Lunevicius	rlunevich@PhD			Foundation Trust	Liverpool				United_Kingdom	School of A University of Liverpool	Liverpool			United_Kingdom
Dr.	Hassan		Magdy	Ab farreha201@MBBCH			Mansoura Faculty of Medicine	Mansoura				Egypt					
Dr.	Mohammed		Magdy	Ab.hoda_734@MBBCH			Aswan Faculty of Medicine	Aswan				Egypt					
Prof.	Reza		Malekzade	malek@tu MD		Aswan University Hospital	Tehran University of Medical Sciences	Tehran				Iran					
Mr.	Kedar	K	Mate	kedar.mat@MSc		Digestive Diseases Research Institute	McGill University	Montreal				Quebec					Canada

Prof. Mr.	Ziad Mubarek	A. Abera	Memish Mengistie	zmemish@MD abmubare@MS		Saudi Ministry of Health Jimma University	Riyadh Jimma		Oromia	Saudi_Arat College of Alfaisal University Ethiopia	Riyadh			Saudi_Arabia		
Dr. Dr. Dr. Dr.	George Felix Farshad Anwar	A. Akpojene	Mensah Ogbo	george.me MD felgbo@ya MPH Pournmalek farshad.po PhD rafay.aku.ε MS	Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute Centre for Health Research	National Institutes of Health Western Sydney University University of British Columbia Contech International Health Consultants	Bethesda Sydney Vancouver Lahore	MD New South Wales British Columbia Punjab		United_States Australia Canada Pakistan	Contech School of Public Health	Lahore	Punjab	Pakistan		
Dr. Prof. Prof. Prof. Prof. Dr.	Mostafa Vafa Saleem Salman Andre Satar		Qorbani Rahimi-Mov Rana Rawaf Renzaho Rezaei	PhD rahimi-mov_rahimi@MD srmrmp@PhD s.rawaf@ir MD andre.renz PhD satar.rezaei PhD	Non-communicable Diseases Research Center Sina Trauma and Surgery Research Center	Albortz University of Medical Sciences Tehran University of Medical Sciences Contech School of Public Health Imperial College London Western Sydney University Kermanshah University of Medical Sciences	Karaj Tehran Lahore London Locked Bag 1797, Penrith 2 Kermanshah	Iran Tehran Punjab		Iran Pakistan United_Kingdom Australia Iran	Contech International Health Consultants	Lahore	Punjab	Pakistan		
Dr. Dr. Dr. Prof. Dr. Dr. Dr.	Mohammad Sadegh Mahdi Mohammad Ali Payman Abdallah M. Juan		Rezaei Mahdi Sahraian Salamati Samy Sanabria	drmsrezaei MD mahdisafdi MD sahraian13 MD psalamati@MD PhD Juan.Sanab MD	Infectious Disease Research Centre with Focus on Nosocomial Infectio Sina Trauma & Surgery Research Center MS Research Center, Neuroscience Institute Sina Trauma and Surgery Research Center J Edwards School of Medicine	Mazandaran University of Medical Sciences Mazandaran University of Medical Sciences Tehran University of Medical Sciences Tehran University of Medical Sciences Ain Shams University Marshall Univeristy	Sari Tehran Tehran Tehran Cairo Huntington	Mazandaran Iran Iran Iran		Iran Iran Iran Iran			United_States	Case Western Reserve University Cleveland	OH	United_States
Dr. Prof. Dr. Dr.	Milena Benn Sadaf Masood	M	Santrici Sepanlou Shaikh	Mil milena.san PhD Sartorius Sepanlou Shaikh	Institute of Social Medicine, Faculty of Medicine Public Health Medicine, School of Nursing and Public Health Digestive Diseases Research Institute	University of Belgrade University of KwaZulu-Natal Tehran University of Medical Sciences Independent Consultant	Belgrade Durban Tehran Karachi	Serbia		Serbia Iran Pakistan	Centre Sch University of Belgrade	Belgrade		Serbia	South_Africa	
Dr. Dr.	Rahman Diego	Augusto Santos	Shiri Silva	rahman.sh PhD diegoaugu PhD	Finnish Institute of Occupational Health, Work Organizations, Work Disability Program, Department of Public Health, Faculty of Medicine	University of Helsinki Federal University of Santa Catarina University of Alabama at Birmingham and Birmingham Veterans Affairs Medical Center	Helsinki Florianopolis	SC - SANTA CATARINA		Finland Brazil						
Dr. Dr. Mr. Dr. Dr.	Jasvinder Abdullah Tenaw Kingsley Tommi	A. Sulieaman Yimer Nnanna	Singh Terkawi Tiruye Ukwajaki Vasankari	jasvinder.n MD asterkawi@MD tenyim09@MPH ukwajaking MD PhD	Department of Anesthesiology Department of Internal Medicine UKK Institute for Health Promotion Research	University of Virginia Debre Markos University Federal Teaching Hospital	Birmingham Charlottesville Debre Markos Abakaliki Tampere	Alabama VA		United_States United_Sta Departmer King Fahad Medical City Ethiopia Nigeria		Riyadh		Saudi_Arat Outcomes Cleveland (Cleveland OH	United_States	
Prof. Mr.	Vasilij Tolassa	Victorovich	Vlassov Wakayo	vlassov@ci MD wakayotoli MS		National Research University Higher School of Economics Jimma University	Moscow Jimma			Russia Ethiopia						
Dr. Mr. Prof. Mr. Dr. Prof.	Andrea Naohiro Mustafa Bassel Aisha Theo		Werdecker Yonemoto Younis Zein Jumaan Vos	andrea.we PhD nyonemot@ MPH younis68@ DrPH b289@geo MS aishaoj@g PhD tvos@uw.ε PhD	Competence Center Mortality-Follow-Up of the German National Cohort Department of Biostatistics, School of Public Health	Federal Institute for Population Research Kyoto University Jackson State University Georgetown University Independent Consultant University of Washington	Wiesbaden Kyoto Jackson Seattle Seattle	Hessen		Germany Japan United_States United_States United_States United_States						
Prof. Prof. Prof. Prof.	Simon Mohsen Christophe Ali	I. L. H.	Hay Murray Mokdad	sihay@uw. DSc naghham@i PhD cjm@uw.ε DPhil mokdaa@i PhD	Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery Institute for Health Metrics and Evaluation Institute for Health Metrics and Evaluation Institute for Health Metrics and Evaluation	University of Oxford University of Washington University of Washington University of Washington	Oxford Seattle Seattle Seattle			United_Kir Institute fo University of Washington United_States United_States United_States	Seattle		Washington	United_States		

Electronic Supplementary Material

Article title:

Burden of obesity in the Eastern Mediterranean Region: findings from the Global Burden of Disease 2015 study

Journal:

International Journal of Public Health

Authors:

GBD 2015 Eastern Mediterranean Region Obesity Collaborators

Corresponding author:

Ali H. Mokdad

Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, United States

Email: mokdaa@uw.edu

E-Table 1: Prevalence of overweight and obesity, with 95% uncertainty intervals (UI) among adults aged 20 years or older (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1980 and 2015)

Country	Socio-demographic Index (SDI)	1980				2015			
		Overweight		Obesity		Overweight		Obesity	
		Males	Females	Males	Females	Males	Females	Males	Females
Afghanistan	Low SDI	29.9 (27.4 – 32.6)	27.3 (24.9 – 29.3)	12.7 (11.0 – 14.6)	17.7 (15.6 – 20.1)	23.1 (20.9 – 25.4)	22.0 (21.1 – 23.0)	7.9 (6.8 – 9.0)	12.1 (11.1 – 13.1)
Bahrain	High-middle SDI	35.8 (33.4 – 37.9)	31.3 (29.7 – 32.6)	18.6 (16.5 – 20.7)	31.0 (28.0 – 34.0)	38.1 (36.7 – 38.9)	32.0 (31.3 – 32.6)	28.1 (25.8 – 30.7)	38.2 (35.3 – 41.2)
Djibouti	Low-middle SDI	9.7 (7.3 – 12.2)	13.2 (11.2 – 15.2)	2.3 (1.3 – 3.8)	3.3 (2.4 – 4.5)	25.3 (22.0 – 28.3)	27.1 (25.5 – 28.5)	9.8 (6.4 – 13.8)	19.4 (15.8 – 23.3)
Egypt	Middle SDI	34.2 (31.3 – 36.9)	33.1 (31.5 – 34.8)	11.2 (9.8 – 12.8)	23.8 (21.3 – 26.4)	41.8 (40.6 – 43.1)	32.4 (32.1 – 32.6)	27.0 (24.7 – 29.3)	43.0 (40.3 – 45.6)
Iran	High-middle SDI	23.9 (21.0 – 26.7)	31.7 (29.4 – 33.8)	5.8 (4.9 – 6.8)	16.6 (14.6 – 18.8)	35.3 (33.0 – 37.5)	32.3 (30.5 – 34.0)	13.8 (12.5 – 15.1)	24.1 (21.9 – 26.2)
Iraq	Middle SDI	36.7 (34.6 – 38.8)	30.7 (29.6 – 32.0)	23.2 (20.9 – 25.5)	33.9 (30.8 – 36.8)	37.2 (36.0 – 38.8)	30.6 (29.7 – 31.2)	25.8 (23.5 – 28.0)	37.0 (34.0 – 39.9)
Jordan	High-middle SDI	36.4 (33.8 – 38.7)	31.4 (29.5 – 32.7)	14.8 (13.0 – 16.6)	27.2 (24.6 – 29.9)	41.6 (40.6 – 42.6)	30.1 (30.0 – 30.3)	31.8 (29.3 – 34.1)	43.4 (40.9 – 45.8)
Kuwait	High SDI	29.6 (27.7 – 31.1)	33.8 (33.0 – 34.7)	23.0 (20.5 – 25.5)	38.9 (36.1 – 41.6)	37.0 (37.1 – 37.0)	32.4 (32.5 – 32.1)	41.4 (39.1 – 43.5)	49.5 (47.8 – 51.4)
Lebanon	High-middle SDI	33.7 (30.7 – 36.8)	30.5 (27.8 – 33.1)	9.2 (8.0 – 10.5)	12.8 (11.1 – 14.6)	42.6 (40.8 – 43.9)	33.1 (31.7 – 34.4)	22.2 (20.2 – 24.1)	24.6 (22.4 – 27.0)
Libya	Middle SDI	38.1 (35.8 – 40.8)	32.2 (30.8 – 33.5)	16.9 (14.8 – 18.9)	27.7 (25.0 – 30.7)	39.6 (37.6 – 41.3)	32.3 (31.2 – 33.1)	21.8 (19.6 – 24.2)	34.9 (31.9 – 38.2)
Morocco	Low-middle SDI	24.74 (22.0 – 27.7)	24.8 (22.7 – 27.2)	6.0 (5.1 – 7.0)	11.4 (9.7 – 13.1)	37.2 (34.3 – 40.0)	32.4 (30.5 – 33.9)	11.8 (10.4 – 13.3)	23.1 (20.6 – 25.7)
Oman	High-middle SDI	27.0 (24.3 – 29.9)	29.0 (26.8 – 30.8)	9.2 (7.9 – 10.5)	20.2 (18.2 – 22.6)	39.6 (38.1 – 40.9)	31.8 (31.0 – 32.4)	24.8 (22.6 – 27.1)	37.9 (35.2 – 40.8)
Pakistan	Low-middle SDI	16.8 (14.1 – 20.0)	21.3 (17.6 – 24.7)	3.6 (2.7 – 4.7)	7.0 (5.5 – 8.8)	24.3 (21.2 – 27.3)	24.4 (22.6 – 26.4)	7.3 (5.9 – 9.1)	14.5 (12.3 – 16.9)
Palestine	Middle SDI	27.0 (24.3 – 29.8)	32.4 (30.5 – 34.6)	10.1 (8.8 – 11.6)	21.0 (18.6 – 23.2)	38.6 (37.5 – 39.6)	37.2 (36.6 – 37.4)	24.5 (22.4 – 26.6)	38.1 (35.6 – 40.5)
Qatar	High-middle SDI	39.1 (39.2 – 39.2)	18.6 (18.1 – 19.4)	43.5 (40.6 – 46.2)	53.0 (49.4 – 56.2)	36.8 (36.7 – 36.8)	23.6 (23.7 – 23.6)	42.5 (40.1 – 44.8)	52.4 (50.3 – 54.6)
Saudi Arabia	High-middle SDI	32.4 (31.3 – 33.2)	32.0 (31.1 – 32.5)	11.4 (11.0 – 12.0)	26.5 (25.7 – 27.5)	38.8 (38.3 – 39.2)	30.3 (30.2 – 30.5)	29.8 (28.9 – 30.7)	43.1 (42.1 – 44.1)
Somalia	Low SDI	9.4 (7.2 – 11.7)	18.0 (15.7 – 20.2)	2.5 (1.4 – 4.2)	8.4 (6.5 – 10.7)	10.7 (8.3 – 13.2)	19.4 (17.6 – 21.1)	2.5 (1.5 – 4.0)	10.5 (8.3 – 12.8)
Sudan	Low-middle SDI	19.7 (17.4 – 22.2)	28.0 (26.0 – 29.9)	5.5 (4.6 – 6.5)	18.2 (16.0 – 20.5)	28.6 (26.1 – 31.1)	30.3 (28.7 – 31.6)	11.5 (10.0 – 13.1)	28.3 (25.6 – 31.2)
Syria	Middle SDI	38.5 (36.1 – 41.1)	34.3 (32.4 – 36.0)	14.2 (12.6 – 15.9)	23.6 (21.3 – 26.0)	43.1 (40.9 – 45.0)	35.1 (33.7 – 36.2)	19.9 (17.9 – 22.0)	30.2 (27.6 – 33.0)
Tunisia	Middle SDI	27.2 (24.5 – 30.5)	29.0 (26.6 – 31.6)	5.4 (4.5 – 6.3)	13.6 (11.8 – 15.5)	40.1 (37.2 – 42.4)	33.4 (31.6 – 35.1)	12.1 (10.7 – 13.8)	23.3 (20.8 – 25.8)
United Arab Emirates	High SDI	38.5	28.9	25.2	31.2	38.1	31.4	25.7	31.6

		(37.2 – 39.9)	(27.6 – 30.1)	(22.6 – 27.5)	(28.1 – 34.1)	(36.4 – 39.8)	(30.3 – 32.5)	(23.2 – 28.2)	(28.9 – 34.6)
Yemen	Low-middle SDI	13.5 (11.6 – 15.6)	19.91 (17.3 – 22.7)	1.9 (1.6 – 2.3)	4.8 (3.9 – 5.8)	24.5 (21.9 – 27.3)	25.4 (23.9 – 26.8)	5.3 (4.5 – 6.3)	12.8 (11.5 – 14.2)

E-Table 2: Prevalence of overweight and obesity, with 95% uncertainty intervals (UI), among children aged 2–19 years (Global Burden of Disease 2015 study, Eastern Mediterranean countries, 1980 and 2015)

Country	Socio-demographic Index (SDI)	1980				2015			
		Overweight		Obesity		Overweight		Obesity	
		Males	Females	Males	Females	Males	Females	Males	Females
Afghanistan	Low SDI	7.9 (5.8–10.2)	14.1 (11.2–17.3)	4.0 (2.8–5.6)	6.2 (4.7–8.2)	4.9 (3.7–6.3)	9.3 (7.2–11.6)	2.1 (1.5–3.0)	4.1 (2.9–5.5)
Bahrain	High-middle SDI	11.2 (8.7–14.1)	15.9 (12.9–19.1)	7.8 (5.6–10.2)	9.3 (7.0–12.2)	12.0 (9.4–14.9)	12.6 (9.9–15.7)	7.7 (5.6–10.4)	7.8 (5.9–10.4)
Djibouti	Low-middle SDI	4.8 (3.1–7.3)	8.8 (6.1–11.4)	2.1 (1.0–3.7)	3.6 (2.1–5.7)	11.6 (8.2–15.6)	15.5 (12.5–19.4)	7.4 (3.9–11.6)	10.5 (6.9–15.1)
Egypt	Middle SDI	10.0 (7.5–12.9)	15.1 (11.8–18.8)	2.8 (1.9–3.8)	4.3 (3.1–5.9)	17.1 (13.6–20.6)	18.3 (15.3–22.0)	9.4 (6.9–12.2)	11.0 (8.4–13.9)
Iran	High-middle SDI	4.1 (3.1–5.4)	6.4 (4.7–8.4)	1.2 (0.8–1.7)	1.7 (1.2–2.3)	8.9 (6.8–11.1)	10.3 (7.7–12.7)	4.3 (3.0–5.7)	4.5 (3.2–6.1)
Iraq	Middle SDI	8.6 (6.6–11.2)	11.4 (8.9–14.4)	3.7 (2.6–5.0)	4.2 (3.0–5.7)	9.9 (7.8–12.4)	12.7 (10.0–15.3)	5.6 (4.1–7.5)	5.8 (4.3–7.7)
Jordan	High-middle SDI	6.3 (4.6–8.3)	9.6 (7.3–12.3)	1.6 (1.1–2.2)	2.3 (1.6–3.2)	13.0 (10.2–16.3)	15.4 (12.7–18.5)	6.3 (4.6–8.2)	6.0 (4.6–7.8)
Kuwait	High SDI	12.2 (9.6–15.3)	15.6 (13.0–18.6)	11.0 (8.1–14.0)	11.0 (8.4–13.9)	19.1 (16.2–21.8)	20.6 (17.5–23.6)	22.1 (17.8–27.0)	19.2 (15.2–23.4)
Lebanon	High-middle SDI	8.1 (6.2–10.4)	10.2 (8.0–12.4)	4.1 (2.9–5.6)	3.7 (2.6–5.0)	13.1 (10.1–15.9)	14.1 (11.4–17.4)	7.5 (5.5–9.7)	6.4 (4.8–8.5)
Libya	Middle SDI	12.8 (9.8–15.6)	17.3 (14.2–20.6)	7.5 (5.5–10.0)	8.7 (6.4–11.2)	13.7 (10.9–16.7)	17.8 (14.4–21.3)	8.8 (6.6–11.4)	9.8 (7.4–12.5)
Morocco	Low-middle SDI	6.4 (4.8–8.5)	10.4 (8.0–13.4)	2.1 (1.5–2.9)	2.6 (1.9–3.6)	10.8 (8.3–13.9)	13.3 (10.4–16.3)	4.7 (3.3–6.5)	4.9 (3.5–6.5)
Oman	High-middle SDI	5.9 (4.3–7.9)	10.7 (8.5–13.2)	2.6 (1.8–3.6)	5.5 (4.0–7.4)	16.2 (12.7–19.5)	16.4 (13.4–19.4)	11.6 (8.6–14.9)	12.0 (9.1–15.3)
Pakistan	Low-middle SDI	5.5 (3.9–7.8)	5.8 (4.1–7.6)	3.0 (2.0–4.5)	3.0 (2.0–4.5)	3.4 (2.3–4.7)	5.3 (3.8–7.7)	2.0 (1.2–2.9)	2.2 (1.4–3.3)
Palestine	Middle SDI	7.0 (5.3–9.3)	9.3 (7.1–11.9)	2.0 (1.4–2.7)	2.3 (1.6–3.2)	11 (8.5–14.1)	12.3 (9.4–15.3)	4.6 (3.3–6.1)	4.9 (3.5–6.5)
Qatar	High-middle SDI	14.7 (12.2–17.1)	9.0 (7.2–10.8)	20.5 (16.1–25.3)	10.6 (7.9–13.8)	16.1 (13.3–18.7)	13.4 (11–16.0)	20.8 (16.6–25.2)	13.5 (10.3–17.2)
Saudi Arabia	High-middle SDI	6.1 (5.5–6.8)	11.1 (10.2–12.0)	2.6 (2.2–3.0)	4.5 (4.0–5.1)	15.3 (14.2–16.5)	19.3 (18.0–20.5)	11.1 (10.1–12.2)	13.8 (12.5–15.2)
Somalia	Low SDI	4.5 (2.8–6.4)	6.3 (4.5–8.1)	2.6 (1.3–4.8)	2.8 (1.7–4.4)	4.3 (2.7–6.2)	7.0 (5.0–9.3)	2.5 (1.2–4.3)	3.2 (2.0–4.8)
Sudan	Low-middle SDI	2.9 (2.1–3.7)	5.4 (4.1–6.7)	1.5 (1.0–2.2)	2.2 (1.5–3.2)	3.3 (2.4–4.4)	5.3 (4.0–6.9)	2.1 (1.5–3.0)	2.7 (2.0–3.6)
Syria	Middle SDI	10.0 (7.6–12.5)	12.3 (9.6–15.2)	3.6 (2.4–4.9)	5.3 (3.8–7.0)	13.3 (10.7–16.4)	13.0 (10.5–16.0)	7.4 (5.4–9.8)	6.3 (4.6–8.3)
Tunisia	Middle SDI	6.0 (4.4–8.0)	7.4 (5.7–9.3)	1.4 (1.0–2.0)	2.1 (1.4–2.8)	13.0 (9.9–16.5)	13.5 (10.7–16.6)	4.1 (2.9–5.5)	4.2 (3.1–5.8)
United Arab Emirates	High SDI	17.2	18.1	10.5	9.7	16.3	17.6	11.3	10.3

		(13.9 – 20.7)	(14.2 – 21.7)	(8.0 – 13.4)	(7.3 – 12.7)	(12.9 – 19.5)	(14.0 – 21.6)	(8.5 – 14.2)	(7.9 – 13.3)
Yemen	Low-middle SDI	2.7 (2.0 – 3.6)	9.3 (7.1 – 11.6)	0.5 (0.3 – 0.6)	2.5 (1.7 – 3.4)	5.1 (3.8 – 6.7)	8.6 (6.6 – 11.0)	1.3 (0.9 – 1.8)	3.7 (2.6 – 5.2)